



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Giant unruptured sinus valsalva aneurysm: an unusual cause of typical angina

Schönrath, Felix ; Alkadhi, Hatem ; Landmesser, Ulf ; Falk, Volkmar ; Plass, André

DOI: <https://doi.org/10.1093/eurheartj/ehs476>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-70922>

Journal Article

Published Version

Originally published at:

Schönrath, Felix; Alkadhi, Hatem; Landmesser, Ulf; Falk, Volkmar; Plass, André (2013). Giant unruptured sinus valsalva aneurysm: an unusual cause of typical angina. *European Heart Journal*, 34(21):1608.

DOI: <https://doi.org/10.1093/eurheartj/ehs476>

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehs476

Online publish-ahead-of-print 12 January 2013

Giant unruptured sinus valsalva aneurysm: an unusual cause of typical angina

Felix Schönmath^{1*}, Hatem Alkadhi², Ulf Landmesser³, Volkmar Falk¹, and André Plass¹


¹Clinic for Cardiovascular Surgery, University Hospital of Zurich, Zurich, Switzerland; ²Clinic for Diagnostic and Interventional Radiology, University Hospital of Zurich, Zurich, Switzerland; and ³Clinic for Cardiology, University Hospital of Zurich, Zurich, Switzerland

* Corresponding author. Tel: +41 44 2553298, Fax: +41 44 2554446. Email: felix.schoenrath@usz.ch

This paper was guest edited by Brahmajee Karthik Nallamothu, University of Michigan, Ann Arbor, MI, USA

A 67-year-old man presented with dyspnoea and angina symptoms in a peripheral hospital. A CT scan of the chest was performed because of suspicion of pulmonary embolism. After application of the contrast agent, an anaphylactic shock occurred. After successful reanimation, a transoesophageal echocardiography displayed a large sinus Valsalva aneurysm (diameter 75 mm). Thereafter, the patient was referred to our hospital.

Magnetic resonance imaging demonstrated a close anatomical relation between the aneurysm and the left coronary artery (Panel B: arrow—sinus Valsalva aneurysm, Panel C: arrow—left main artery, asterisk—sinus Valsalva aneurysm). A coronary angiogram was performed and revealed occlusion of the left main artery caused by the large unruptured aneurysm. The left coronary system filled partly retrograde (Panels D and E, Supplementary material online, Videos S2 and S3). Echocardiography during open heart surgery showed the aneurysm to be filled with a mobile thrombus (Panel A, Supplementary material online, Video S1). The intra-operative assessment of the aneurysm showed the origin inferior to the left main coronary artery continuing laterally just posterior to the pulmonary artery (Panel F, black arrow: aneurysm, white arrow and Panel G—thrombus). Because the aneurysm adhered to the left coronary ostium, coronary bypass grafts to the left anterior descending artery and circumflex artery were performed using the left (LIMA) and right (as T-Graft from LIMA) mammary arteries. The aneurysm was resected and the aortic root was replaced with a biological composite graft. The patient had a post-operative course without complication and a fast recovery.



Supplementary material is available at *European Heart Journal* online.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2013. For permissions please email: journals.permissions@oup.com